
NUMBER OF UNMARKED OLDER JUVENILE CHINOOK MEASURED IN THE SACRAMENTO RIVER



DWR-DES 22 DECEMBER 2014


Preliminary data from DFW, FWS, GCID, and CDEC; subject to revision.

^{*}Older juvenile Chinook defined as all Chinook greater than or equal to the minimum winter run length-at-date criteria and less than the maximum size included in the length-at-date criteria (Frank Fisher model) for which a race is assigned on a given sampling date.

^{**}Turbidity is a discrete measurement and is not measured continuously. Therefore, data are interpolated on days when turbidity was not measured unless data are missing for more than five days.

^{***}Trap was pulled on 10/28/14 due to extremely turbid conditions, heavy debris, and high number of listed winter run Chinook and has resumed since 11/5/14.

NUMBER OF UNMARKED OLDER JUVENILE CHINOOK MEASURED IN THE LOWER SACRAMENTO RIVER AND CHIPPS ISLAND




DWR-DES 22 DECEMBER 2014

Preliminary data from DFW, FWS, and CDEC; subject to revision.
*Older juvenile Chinook defined as all Chinook greater than or equal to the minimum winter run length-at-date criteria and less than the maximum size included in the length-at-date criteria (Frank Fisher Model) for which a race is assigned on a given sampling date.

^{**}Turbidity is a discrete measurement and is not measured continuously. Therefore, data are interpolated on days when turbidity was not measured unless data are missing for more than five days. Knights Landing turbidity measured in FTU, which should be roughly equivalent to NTU.

^{***}Sacramento area seine route consists of the following seine sites: Verona, Elkhorn, Sand Cove, Discovery Park, American River, Miller Park, Sherwood Harbor, and Garcia Bend. Bars are stacked if Chinook caught from the trawl and seines are from the same day.

NUMBER OF UNMARKED OLDER JUVENILE CHINOOK AND STEELHEAD MEASURED IN THE SAN JOAQUIN RIVER




DWR-DES 22 DECEMBER 2014
Preliminary data from FWS and CDEC; subject to revision.

^{*}Older juvenile Chinook defined as all Chinook greater than or equal to the minimum winter run length-at-date criteria and less than the maximum size included in the length-at-date criteria (Frank Fisher model) for which a race is assigned on a given sampling date.

^{**}Turbidity is a discrete measurement and is not measured continuously. Therefore, data are interpolated on days when turbidity was not measured unless data are missing for more than five days.

NUMBER OF UNMARKED FRY/SMOLT CHINOOK **MEASURED IN THE SACRAMENTO RIVER**




DWR-DES 22 DECEMBER 2014

Preliminary data from DFW, FWS, GCID, and CDEC; subject to revision.
*Fry/smolt Chinook defined as all Chinook less than the minimum winter run length-at-date criteria (Frank Fisher model).

^{**}Turbidity is a discrete measurement and is not measured continuously. Therefore, data are interpolated on days when turbidity was not measured unless data are missing for more than five days.

^{***}Trap was pulled on 10/28/14 due to extremely turbid conditions, heavy debris, and high number of listed winter run Chinook and has resumed since 11/5/14.

NUMBER OF UNMARKED FRY/SMOLT CHINOOK MEASURED IN THE LOWER SACRAMENTO RIVER AND CHIPPS ISLAND



DWR-DES 22 DECEMBER 2014


Preliminary data from DFW, FWS, and CDEC; subject to revision.

*Fry/smolt Chinook defined as all Chinook less than the minimum winter run length-at-date criteria (Frank Fisher model).

^{**}Turbidity is a discrete measurement and is not measured continuously. Therefore, data are interpolated on days when turbidity was not measured unless data are missing for more than five days. Knights Landing turbidity measured in FTU, which should be roughly equivalent to NTU.

^{***}Sacramento area seine route consists of the following seine sites: Verona, Elkhorn, Sand Cove, Discovery Park, American River, Miller Park, Sherwood Harbor, and Garcia Bend. Bars are stacked if Chinook caught from the trawl and seines are from the same day.


NUMBER OF UNMARKED FRY/SMOLT CHINOOK **MEASURED IN THE SAN JOAQUIN RIVER**



Preliminary data from FWS and CDEC; subject to revision.
*Fry/smolt Chinook defined as all Chinook less than the minimum winter run length-at-date criteria (Frank Fisher model).

^{**}Turbidity is a discrete measurement and is not measured continuously. Therefore, data are interpolated on days when turbidity was not measured unless data are missing for more than five days.

WATER TEMPERATURE AND FLOW MEASURED AT MILL AND DEER CREEK



Data Acquisition:

All data are preliminary and subject to revision.

The estimated passage data for the Red Bluff Diversion Dam were obtained directly from the US Fish and Wildlife Service (FWS), Red Bluff Fish and Wildlife Office (http://www.fws.gov/redbluff/rbdd biweekly.aspx).

The catch data for Glenn-Colusa Irrigation District (GCID) were obtained directly from GCID.

The catch data for Tisdale Weir and Knights Landing were obtained directly from the California Department of Fish and Wildlife (DFW).¹, North Central Region.

Sacramento River Trawl, Sacramento Area Beach Seine, and Chipps Island Trawl data were obtained directly from FWS, Stockton Fish and Wildlife Office (http://www.fws.gov/stockton/jfmp/).

Mossdale Trawl data were either obtained directly from FWS, Stockton Fish and Wildlife Office or from DFW (Region 4).

The hydrology data were either downloaded from the California Data Exchange Center (CDEC) (http://cdec.water.ca.gov) or obtained directly from the California Department of Water Resources, Operations Control Office.

_

¹ Formerly known as the California Department of Fish and Game (DFG).